


# Regeneration (biology)

From Wikipedia, the free encyclopedia


In biology, **regeneration** is the process of renewal, restoration, and growth that makes genomes, cells, organs, organisms, and ecosystems resilient to natural fluctuations or events that cause disturbance or damage. Every species is capable of regeneration, from bacteria to humans.<sup>[1][2]</sup> At its most elementary level, regeneration is mediated by the molecular processes of DNA synthesis.<sup>[3][4]</sup> Regeneration in biology, however, mainly refers to the morphogenic processes that characterizes the phenotypic plasticity of traits allowing multi-cellular organisms to repair and maintain the integrity of their physiological and morphological states. Above the genetic level, regeneration is fundamentally regulated by asexual cellular processes.<sup>[5]</sup>

The hydra and the planarian flatworm have long served as model organisms for their highly adaptive regenerative capabilities.<sup>[6]</sup> Once wounded, their cells become activated and start to remodel tissues and organs back to the pre-existing state.<sup>[7]</sup> The urodele (salamander), an amphibian, is possibly the most adept vertebrate order for their capability of regenerating limbs, tails, jaws, eyes and a variety of internal structures.<sup>[1]</sup> The regeneration of organs is a common and widespread adaptive capability among metazoan creatures.<sup>[6]</sup> In a related

context, some animals are able to reproduce asexually through fragmentation, budding, or fission.<sup>[5]</sup> A planarian parent, for example, will constrict, split in the middle, and each half generates a new end to form two clones of the original.<sup>[8]</sup> Echinoderms, such as the starfish, crayfish, many reptiles, and amphibians exhibit remarkable examples of tissue regeneration. The case of autotomy, for example, serves as a defensive function as the animal detaches a limb or tail to avoid capture. After the limb or tail has been autotomized, cells move into action and tissues regenerate.<sup>[9][10][11]</sup> Ecosystems are regenerative as well. Following a disturbance, such as a fire or pest outbreak in a forest, pioneering species will occupy, compete for space, and establish themselves in the newly opened habitat. The new growth of seedlings and community assembly process is known as regeneration in ecology.<sup>[12][13]</sup>



Sun flower sea star regenerates its arms



Dwarf yellow-headed gecko with regenerating tail

## Contents

- 1 Cellular molecular fundamentals
- 2 Tissues and organs
- 3 Model organisms in Bilateria
  - 3.1 Planaria (Platyhelminthes)

- 3.2 Vertebrates
  - 3.2.1 Amphibians
  - 3.2.2 Mice
  - 3.2.3 Humans
    - 3.2.3.1 Fingertips
    - 3.2.3.2 Ribs
    - 3.2.3.3 Liver
    - 3.2.3.4 Kidney
    - 3.2.3.5 Heart
- 4 Notes
- 5 Sources
- 6 External links

## Cellular molecular fundamentals

Pattern formation in the morphogenesis of an animal is regulated by genetic induction factors that put cells to work after damage has occurred. Neural cells, for example, express growth-associated proteins, such as GAP-43, tubulin, actin, an array of novel neuropeptides, and cytokines that induce a cellular physiological response to regenerate from the damage.<sup>[14]</sup> Many of the genes that are involved in the original development of tissues are reinitialized during the regenerative process. Cells in the primordia of zebrafish fins, for example, express four genes from the homeobox *msx* family during development and regeneration.<sup>[15]</sup>

## Tissues and organs

"Regenerative strategies include the rearrangement of pre-existing tissue, the use of adult somatic stem cells and the dedifferentiation and/or transdifferentiation of cells, and more than one mode can operate in different tissues of the same animal. All these strategies result in the re-establishment of appropriate tissue polarity, structure and form."<sup>[16]:873</sup> During the developmental process genes are activated that serve modify the properties of cell as they differentiate into different tissues. Development and regeneration involves the coordination and organization of populations cells into a blastema, which is "a mound of stem cells from which regeneration begins."<sup>[17]</sup> The dedifferentiation of cells means that they lose their tissue specific characteristics as tissues remodel during the regeneration process. The transdifferentiation of cells means that they lose their tissue specific characteristics during the regeneration process and then re-differentiate to a different kind of cell.<sup>[16]</sup>

## Model organisms in Bilateria

### Planaria (Platyhelminthes)

Planaria exhibit an extraordinary ability to regenerate lost body parts. For example, a planarian split lengthwise or crosswise will regenerate into two separate individuals. In one experiment, T. H. Morgan found that a piece corresponding to 1/279th of a planarian could successfully regenerate into a new worm. This size (about 10,000 cells) is typically accepted as the smallest fragment that can regrow into a new planarian. Regeneration of planaria is epimorphic regeneration. After amputation, stump cells form blastema.

## Vertebrates

### Amphibians

Limb regeneration in newts occurs in two major steps, first de-differentiation of adult cells into a stem cell state similar to embryonic cells and second, development of these cells into new tissue more or less the same way it developed the first time.<sup>[18]</sup> Simpler animals like planarian have an enhanced capacity to regenerate because the adults retain clusters of stem cells (neoblast) within their bodies which migrate to the parts that need healing. They then divide and differentiate to grow the missing tissue and organs back.

In salamanders, the regeneration process begins immediately after amputation. Limb regeneration in the axolotl and newt have been extensively studied. After amputation, the epidermis migrates to cover the stump in less than 12 hours, forming a structure called the apical epidermal cap (AEC). Over the next several days there are changes in the underlying stump tissues that result in the formation of a blastema (a mass of dedifferentiated proliferating cells). As the blastema forms, pattern formation genes – such as HoxA and HoxD – are activated as they were when the limb was formed in the embryo.<sup>[19][20]</sup> The distal tip of the limb (the autopod, which is the hand or foot) is formed first in the blastema. The intermediate portions of the pattern are filled in during growth of the blastema by the process of intercalation.<sup>[18][19]</sup> Motor neurons, muscle, and blood vessels grow with the regenerated limb, and reestablish the connections that were present prior to amputation. The time that this entire process takes varies according to the age of the animal, ranging from about a month to around three months in the adult and then the limb becomes fully functional.

In spite of the historically few researchers studying limb regeneration, remarkable progress has been made recently in establishing the neotenous amphibian the axolotl (*Ambystoma mexicanum*) as a model genetic organism. This progress has been facilitated by advances in genomics, bioinformatics, and somatic cell transgenesis in other fields, that have created the opportunity to investigate the mechanisms of important biological properties, such as limb regeneration, in the axolotl.<sup>[21]</sup> The Ambystoma Genetic Stock Center (AGSC) is a self-sustaining, breeding colony of the axolotl supported by the National Science Foundation as a Living Stock Collection. Located at the University of Kentucky, the AGSC is dedicated to supplying genetically well-characterized axolotl embryos, larvae, and adults to laboratories throughout the United States and abroad. An NIH-funded NCR grant has led to the establishment of the Ambystoma EST database, the Salamander Genome Project (SGP) that has led to the creation of the first amphibian gene map and several annotated molecular data bases, and the creation of the research community web portal.<sup>[22]</sup>

### Mice

The mechanism for regeneration in MRL mice has been found and it is related to the deactivation of the p21 gene.<sup>[23][24]</sup>

Adult mammals have limited regenerative capacity compared to most vertebrate embryos/larvae, adult salamanders and fish. The MRL mouse is a strain of mouse that exhibits remarkable regenerative abilities for a mammal. Study of the regenerative process in these animals is aimed at discovering how to duplicate them in humans.

By comparing the differential gene expression of scarless healing MRL mice and poor healing C57BL/6 mice strain, 36 genes have been identified that are good candidates for studying how the healing process differs in MRL mice and other mice.<sup>[25][26]</sup>

The regenerative abilities of MRL mice does not, however, protect them against myocardial infarction, as heart regeneration in adult mammals (neocardiogenesis) is limited because heart muscle cells are nearly all terminally differentiated. MRL mice show the same amount of cardiac injury and scar formation as normal mice after a heart attack.<sup>[27]</sup> Though recent studies provide evidence that this may not be the case, and that MRL mice do regenerate from heart damage. [2] ([http://www.eurekalert.org/pub\\_releases/2001-08/wi-rit080201.php](http://www.eurekalert.org/pub_releases/2001-08/wi-rit080201.php))

## Humans

### Fingertips

Studies in the 1970s showed that children up to the age of 10 or so who lose fingertips in accidents can regrow the tip of the digit within a month provided their wounds are not sealed up with flaps of skin – the de facto treatment in such emergencies. They normally won't have a finger print, and if there is any piece of the finger nail left it will grow back as well, usually in a square shape rather than round.<sup>[28][29]</sup>

In August 2005, Lee Spievack, then in his early sixties, accidentally sliced off the tip of his right middle finger just above the first phalanx. His brother, Dr. Alan Spievack, was researching regeneration and provided him with powdered extracellular matrix, developed by Dr. Stephen Badylak of the McGowan Institute (<http://www.mirm.pitt.edu/>) of Regenerative Medicine. Mr. Spievack covered the wound with the powder, and the tip of his finger re-grew in four weeks.<sup>[30]</sup> The news was released in 2007. Lee Spievack is the first documented case of an adult human regenerating fingertips;<sup>[28]</sup> however, Ben Goldacre has described this as "the missing finger that never was", claiming that fingertips regrow and quoted Simon Kay, professor of hand surgery at the University of Leeds, who from the picture provided by Goldacre described the case as seemingly "an ordinary fingertip injury with quite unremarkable healing"<sup>[31]</sup>

A similar story was reported by CNN. A woman named Deepa Kulkarni lost the tip of her little finger and was initially told by doctors that nothing could be done. Her personal research and consultation with several specialists including Badylak eventually resulted in her undergoing regenerative therapy and regaining her fingertip.<sup>[32]</sup>

### Ribs

There have appeared claims that human ribs could regenerate if the periosteum, the membrane surrounding the rib, were left intact. In one study rib material was used for skull reconstruction and all 12 patients had complete regeneration of the resected rib.<sup>[33]</sup>

### Liver

The human liver is one of the few glands in the body that has the ability to regenerate from as little as 25% of its tissue.<sup>[34]</sup> This is largely due to the unipotency of hepatocytes.<sup>[35]</sup> Resection of liver can induce the proliferation of the remained hepatocytes until the lost mass is restored, where the intensity of the liver's response is directly proportional to the mass resected. For almost 80 years surgical resection of the liver in rodents has been a very useful model to the study of cell proliferation.<sup>[36][37]</sup>

## Kidney

Regenerative capacity of the kidney remains largely unexplored. The basic functional and structural unit of the kidney is nephron, which is mainly composed of four components: the glomerulus, tubules, the collecting duct and peritubular capillaries. The regenerative capacity of the mammalian kidney is limited compared to that of lower vertebrates.

In the mammalian kidney, the regeneration of the tubular component following an acute injury is well known. Recently regeneration of the glomerulus has also been documented. Following an acute injury, the proximal tubule is damaged more, and the injured epithelial cells slough off the basement membrane of the nephron. The surviving epithelial cells, however, undergo migration, dedifferentiation, proliferation, and redifferentiation to replenish the epithelial lining of the proximal tubule after injury. Recently, the presence and participation of kidney stem cells in the tubular regeneration has been shown. However, the concept of kidney stem cells is currently emerging. In addition to the surviving tubular epithelial cells and kidney stem cells, the bone marrow stem cells have also been shown to participate in regeneration of the proximal tubule, however, the mechanisms remain controversial. Recently, studies examining the capacity of bone marrow stem cells to differentiate into renal cells are emerging.<sup>[38]</sup>

Like other organs, the kidney is also known to regenerate completely in lower vertebrates such as fish. Some of the known fish that show remarkable capacity of kidney regeneration are goldfish, skates, rays, and sharks. In these fish, the entire nephron regenerates following injury or partial removal of the kidney.

## Heart

Several animals can regenerate heart damage, but in mammals cardiomyocytes cannot proliferate and heart damage causes scarring and fibrosis.

The long held view was that mammalian cardiomyocytes are terminally differentiated and cannot divide. However inhibition of p38 MAP kinase was found to induce mitosis in adult mammalian cardiomyocytes.<sup>[39]</sup> Treatment with FGF1 and p38 MAP kinase inhibitors regenerates the heart, reduces scarring, and improves cardiac function in rats with cardiac injury.<sup>[40]</sup>

## Notes

- <sup>a b</sup> Carlson, B. M. (2007). *Principles of Regenerative Biology* (<http://www.sciencedirect.com/science/book/9780123694393>) . Elsevier Inc.. pp. 400. ISBN 978-0-12-369439-3. <http://www.sciencedirect.com/science/book/9780123694393>.
- <sup>a</sup> Gabor, M. H.; Hotchkiss, R. D. (1979). "Parameters governing bacterial regeneration and genetic recombination after fusion of *Bacillus subtilis* protoplasts." (<http://jb.asm.org/cgi/content/abstract/137/3/1346>) . *Journal of Bacteriology* **137** (3): 1346–1353. PMC 218319 (<http://www.pubmedcentral.gov/articlerender.fcgi?tool=pmcentrez&artid=218319>) . PMID 108246 (<http://www.ncbi.nlm.nih.gov/pubmed/108246>) . <http://jb.asm.org/cgi/content/abstract/137/3/1346>.
- <sup>a</sup> Himeno, Y.; Engelman, R. W.; Good, R. A. (1992). "Influence of calorie restriction on oncogene expression and DNA synthesis during liver regeneration." (<http://www.pnas.org/content/89/12/5497.abstract>) . *Proceedings of the National Academy of Sciences of the United States of America* **89** (12): 5497–5501. doi:10.1073/pnas.89.12.5497 (<http://dx.doi.org/10.1073/pnas.89.12.5497>) . PMC 49319 (<http://www.pubmedcentral.gov/articlerender.fcgi?tool=pmcentrez&artid=49319>) . PMID 1608960 (<http://www.ncbi.nlm.nih.gov/pubmed/1608960>) . <http://www.pnas.org/content/89/12/5497.abstract>.
- <sup>a</sup> Bryant, P. J.; Fraser, S. E. (1988). "Wound healing, cell communication, and DNA synthesis during imaginal disc regeneration in *Drosophila*." (<http://linkinghub.elsevier.com/retrieve/pii/0012160688902011>) . *Developmental Biology*

- 127 (1): 197–208. doi:10.1016/0012-1606(88)90201-1 (<http://dx.doi.org/10.1016%2F0012-1606%2888%2990201-1>) . PMID 2452103 (<http://www.ncbi.nlm.nih.gov/pubmed/2452103>) .  
<http://linkinghub.elsevier.com/retrieve/pii/0012160688902011>.
5. <sup>a b</sup> Brookes, J. P.; Kumar, A.. "Comparative Aspects of Animal Regeneration." (<http://www.annualreviews.org/doi/full/10.1146/annurev.cellbio.24.110707.175336?select23=Choose>) . *Annu. Rev. Cell Dev. Biol.* **28**: 525–549. doi:10.1146/annurev.cellbio.24.110707.175336?select23=Choose (<http://dx.doi.org/10.1146%2Fannurev.cellbio.24.110707.175336%3Fselect23%3DChoose>) .  
<http://www.annualreviews.org/doi/full/10.1146/annurev.cellbio.24.110707.175336?select23=Choose>.
6. <sup>a b</sup> Sánchez, A. A. (2000). "Regeneration in the metazoans: why does it happen?" (<http://planaria.neuro.utah.edu/publications/BioEssays.pdf>) . *Bioessays* **22** (6): 578–590. doi:10.1002/(SICI)1521-1878(200006)22:6<578::AID-BIES11>3.0.CO;2-# (<http://dx.doi.org/10.1002%2F%28SICI%291521-1878%28200006%2922%3A6%3C578%3A%3AAID-BIES11%3E3.0.CO%3B2-%23>) . PMID 10842312 (<http://www.ncbi.nlm.nih.gov/pubmed/10842312>) .  
<http://planaria.neuro.utah.edu/publications/BioEssays.pdf>.
7. <sup>a</sup> Reddien, P. W.; Alvarado, A. S. (2004). "Fundamentals of planarian regenerations" (<http://www.annualreviews.org/doi/abs/10.1146/annurev.cellbio.20.010403.095114>) . *Annual Review of Cell and Developmental Biology* **20**: 725–757. doi:10.1146/annurev.cellbio.20.010403.095114 (<http://dx.doi.org/10.1146%2Fannurev.cellbio.20.010403.095114>) . PMID 15473858 (<http://www.ncbi.nlm.nih.gov/pubmed/15473858>) .  
<http://www.annualreviews.org/doi/abs/10.1146/annurev.cellbio.20.010403.095114>.
8. <sup>a</sup> Campbell, N. A.. *Biology* (4th ed.). California: The Benjamin Cummings Publishing Company, Inc.. pp. 1206.
9. <sup>a</sup> Wilkie, I. (2001). "Autotomy as a prelude to regeneration in echinoderms." (<http://onlinelibrary.wiley.com/doi/10.1002/jemt.1185/abstract>) . *Microscopy Research and Technique* **55** (6): 369–396. doi:10.1002/jemt.1185 (<http://dx.doi.org/10.1002%2Fjemt.1185>) . PMID 11782069 (<http://www.ncbi.nlm.nih.gov/pubmed/11782069>) .  
<http://onlinelibrary.wiley.com/doi/10.1002/jemt.1185/abstract>.
10. <sup>a</sup> Maiorana, V. C. (1977). "Tail autotomy, functional conflicts and their resolution by a salamander." . *Nature* **226** (5594): 533–535. doi:10.1038/265533a0 (<http://dx.doi.org/10.1038%2F265533a0>) .
11. <sup>a</sup> Maginnis, T. L. (2006). "The costs of autotomy and regeneration in animals: a review and framework for future research." (<http://beheco.oxfordjournals.org/content/17/5/857.full>) . *Behavioural Ecology* **7** (5): 857–872. doi:10.1093/beheco/arl010 (<http://dx.doi.org/10.1093%2Fbeheco%2Farl010>) .  
<http://beheco.oxfordjournals.org/content/17/5/857.full>.
12. <sup>a</sup> Dietze, M. C.; Clark, J. S. (2008). "Changing the gap dynamics paradigm: Vegetative regenerative control on forest response to disturbance." (<http://coweeta.uga.edu/publications/10300.pdf>) . *Ecological Monographs* **78** (3): 331–347. doi:10.1890/07-0271.1 (<http://dx.doi.org/10.1890%2F07-0271.1>) .  
<http://coweeta.uga.edu/publications/10300.pdf>.
13. <sup>a</sup> Bailey, J. D.; Covington, W. W. (2002). "Evaluation ponderosa pine regeneration rates following ecological restoration treatments in northern Arizona, USA." (<http://library.eri.nau.edu/gsdll/collect/erilibra/index/assoc/HASH0e9a.dir/doc.pdf>) . *Forest Ecology and Management* **155**: 271–278. doi:10.1016/S0378-1127(01)00564-3 (<http://dx.doi.org/10.1016%2FS0378-1127%2801%2900564-3>) .  
<http://library.eri.nau.edu/gsdll/collect/erilibra/index/assoc/HASH0e9a.dir/doc.pdf>.
14. <sup>a</sup> Fu, S. Y.; Gordon, T. (1997). "The cellular and molecular basis of peripheral nerve regeneration." (<http://www.springerlink.com/content/675770336wq42078/>) . *Molecular Neurobiology* **14** (1-2): 67–116. doi:10.1007/BF02740621 (<http://dx.doi.org/10.1007%2F02740621>) . PMID 9170101 (<http://www.ncbi.nlm.nih.gov/pubmed/9170101>) .  
<http://www.springerlink.com/content/675770336wq42078/>.
15. <sup>a</sup> Akimenko, M.; Johnson, S. L.; Wsoterfield, M.; Ekker, M. (1996). "Differential induction of four *msx* homeobox genes during fin development and regeneration in zebrafish." (<http://dev.biologists.org/content/121/2/347.full.pdf>) . *Development* **121** (2): 347–357. PMID 7768177 (<http://www.ncbi.nlm.nih.gov/pubmed/7768177>) .  
<http://dev.biologists.org/content/121/2/347.full.pdf>.
16. <sup>a b</sup> Alvarado, A. S.; Tsonis, P. A. (2006). "Bridging the regeneration gap: genetic insights from diverse animal models." (<http://planaria.neuro.utah.edu/publications/NRG2006.pdf>) . *Nat. Rev. Genet.* **7** (11): 873–884. doi:10.1038/nrg1923 (<http://dx.doi.org/10.1038%2Fnrg1923>) . PMID 17047686 (<http://www.ncbi.nlm.nih.gov/pubmed/17047686>) .  
<http://planaria.neuro.utah.edu/publications/NRG2006.pdf>.

17. ^ Kumar, A.; Godwin, J. W.; Gates, P. B.; Garza-Garcia, A. A.; Brokes, J. P. (2007). "Molecular Basis for the Nerve Dependence of Limb Regeneration in an Adult Vertebrate." (<http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pmcentrez&artid=2696928>). *Science* **318** (5851): 772. doi:10.1126/science.1147710 (<http://dx.doi.org/10.1126%2Fscience.1147710>). PMC 2696928 (<http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pmcentrez&artid=2696928>). PMID 17975060 (<http://www.ncbi.nlm.nih.gov/pubmed/17975060>). <http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pmcentrez&artid=2696928>.
18. ^ <sup>a</sup> <sup>b</sup> Odelberg SJ. Unraveling the molecular basis for regenerative cellular plasticity. *PLoS Biol.* 2004 Aug;2(8):E232. PMID 15314652
19. ^ <sup>a</sup> <sup>b</sup> Bryant, S.V., Endo, T. and Gardiner, D.M. Vertebrate limb regeneration and the origin of limb stem cells. *Int. J. Dev. Biol.* 2002 46:887-896. PMID 12455626
20. ^ Mullen LM, Bryant SV, Torok MA, Blumberg B, Gardiner DM. Nerve dependency of regeneration: the role of Distal-less and FGF signaling in amphibian limb regeneration. *Development.* 1996 Nov;122 (11):3487-97. PMID 8951064
21. ^ Endo T, Bryant SV, Gardiner DM. A stepwise model system for limb regeneration. *Dev Biol.* 2004 Jun 1;270(1):135-45. PMID 15136146
22. ^ <http://www.ambystoma.org>
23. ^ Bedelbaeva K, Snyder A, Gourevitch D, Clark L, Zhang X-M, Leferovich J, Cheverud JM, Lieberman P, Heber-Katz E (March 2010). "Lack of p21 expression links cell cycle control and appendage regeneration in mice" (<http://www.pnas.org/content/early/2010/03/08/1000830107.abstract>). *Proceedings of the National Academy of Sciences* **107** (11): 5845–50. doi:10.1073/pnas.1000830107 (<http://dx.doi.org/10.1073%2Fpnas.1000830107>). PMC 2851923 (<http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pmcentrez&artid=2851923>). PMID 20231440 (<http://www.ncbi.nlm.nih.gov/pubmed/20231440>). <http://www.pnas.org/content/early/2010/03/08/1000830107.abstract>. Lay summary (<http://www.physorg.com/news187879295.html>) – *PhysOrg.com*.
24. ^ Humans Could Regenerate Tissue Like Newts By Switching Off a Single Gene (<http://www.popsci.com/science/article/2010-03/humans-could-regenerate-tissue-newts-switchin-single-gene>)
25. ^ *Biochem Biophys Res Commun.* 2005 Apr 29;330(1):117-22. PMID 15781240
26. ^ Mansuo L, Hayashi, B. S. Shankaranarayana Rao, Jin-Soo Seo, Han-Saem Choi, Bridget M. Dolan, Se-Young Choi, Sumantra Chattarji, and Susumu Tonegawa (2007 July). "Inhibition of p21-activated kinase rescues symptoms of fragile X syndrome in mice" (<http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pmcentrez&artid=1899186>). *Proceedings of the National Academy of Sciences* **104** (27): 11489. doi:10.1073/pnas.0705003104 (<http://dx.doi.org/10.1073%2Fpnas.0705003104>). PMC 1899186 (<http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pmcentrez&artid=1899186>). PMID 17592139 (<http://www.ncbi.nlm.nih.gov/pubmed/17592139>). <http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pmcentrez&artid=1899186>.
27. ^ Abdullah I, Lepore JJ, Epstein JA, Parmacek MS, Gruber PJ (2005 Mar-April). "MRL mice fail to heal the heart in response to ischemia-reperfusion injury." *Wound Repair Regen* **13** (2): 205–208. doi:10.1111/j.1067-1927.2005.130212.x (<http://dx.doi.org/10.1111%2Fj.1067-1927.2005.130212.x>). PMID 15828946 (<http://www.ncbi.nlm.nih.gov/pubmed/15828946>).
28. ^ <sup>a</sup> <sup>b</sup> Weintraub, Arlene (MAY 24, 2004). "The Geniuses Of Regeneration" ([http://www.businessweek.com/magazine/content/04\\_21/b3884008\\_mz001.htm](http://www.businessweek.com/magazine/content/04_21/b3884008_mz001.htm)). *BusinessWeek*. [http://www.businessweek.com/magazine/content/04\\_21/b3884008\\_mz001.htm](http://www.businessweek.com/magazine/content/04_21/b3884008_mz001.htm).
29. ^ Illingworth, Cynthia M. 1974. Trapped fingers and amputated fingertips in children. *J. Ped. Surgery* 9:853-858.
30. ^ "Regeneration recipe: Pinch of pig, cell of lizard" (<http://www.msnbc.msn.com/id/17171083/>). Associated Press. MSNBC. February 19, 2007. <http://www.msnbc.msn.com/id/17171083/>. Retrieved October 24, 2008.
31. ^ Goldacre, Ben (May 3, 2008). "The missing finger that never was" (<http://www.guardian.co.uk/science/2008/may/03/medicalresearch.health>). *The Guardian*. <http://www.guardian.co.uk/science/2008/may/03/medicalresearch.health>.
32. ^ Woman's persistence pays off in regenerated fingertip (<http://www.cnn.com/2010/HEALTH/09/09/pinky.regeneration.surgery/index.html>) by Elizabeth Cohen. CNN website, September 9, 2010 4:51 p.m., page found 2010-09-16.
33. ^ Munro IR, Guyuron B (November 1981). "Split-Rib Cranioplasty". *Annals of Plastic Surgery* **7** (5): 341–346. doi:10.1097/00000637-198111000-00001 (<http://dx.doi.org/10.1097%2F00000637-198111000-00001>). PMID 7332200 (<http://www.ncbi.nlm.nih.gov/pubmed/7332200>).



34. ^ "Liver Regeneration Unplugged" (<http://www.bio-medicine.org/medicine-news/Liver-Regeneration-Unplugged-19988-1/>) . Bio-Medicine. 2007-04-17. <http://www.bio-medicine.org/medicine-news/Liver-Regeneration-Unplugged-19988-1/>. Retrieved 2007-04-17.
35. ^ Michael, Dr. Sandra Rose (2007). "Bio-Scalar Technology: Regeneration and Optimization of the Body-Mind Homeostasis" ([http://eesystem.com/docs/AAAAM%202007%20long%20biography%20abstr\\_.pdf](http://eesystem.com/docs/AAAAM%202007%20long%20biography%20abstr_.pdf)) (PDF). *15th Annual AAAAM Conference*; 2. [http://eesystem.com/docs/AAAAM%202007%20long%20biography%20abstr\\_.pdf](http://eesystem.com/docs/AAAAM%202007%20long%20biography%20abstr_.pdf). Retrieved October 24, 2008.
36. ^ Higgins, GM and RM Anderson RM (1931). "Experimental pathology of the liver. I. Restoration of the liver of the white rat following partial surgical removal". *Arch. Pathol.* **12**: 186–202.
37. ^ Michalopoulos, GK and MC DeFrances (April 4, 1997). "Liver regeneration". *Science* **276** (5309): 60–66. doi:10.1126/science.276.5309.60 (<http://dx.doi.org/10.1126/science.276.5309.60>) . PMID 9082986 (<http://www.ncbi.nlm.nih.gov/pubmed/9082986>) .
38. ^ Kurinji Singaravelu et al.(July 2009). "In Vitro Differentiation of MSC into Cells with a Renal Tubular Epithelial-Like Phenotype". *Renal Failure* **31**(6):492-502. <http://www.informaworld.com/smpp/content~content=a913452182~db=all~jumpType=rss>
39. ^ [1] (<http://genesdev.cshlp.org/content/19/10/1175.full>)
40. ^ Felix B. Engel, Patrick C. H. Hsieh, Richard T. Lee, Mark T. Keating (2006 October). "FGF1/p38 MAP kinase inhibitor therapy induces cardiomyocyte mitosis, reduces scarring, and rescues function after myocardial infarction" (<http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pmcentrez&artid=1622860>) . *Proceedings of the National Academy of Sciences* **103** (42): 15546–15551. doi:10.1073/pnas.0607382103 (<http://dx.doi.org/10.1073/pnas.0607382103>) . PMC 1622860 (<http://www.pubmedcentral.gov/articlerender.fcgi?tool=pmcentrez&artid=1622860>) . PMID 17032753 (<http://www.ncbi.nlm.nih.gov/pubmed/17032753>) . <http://www.pubmedcentral.nih.gov/articlerender.fcgi?tool=pmcentrez&artid=1622860>.

## Sources

1. Tanaka EM. Cell differentiation and cell fate during urodele tail and limb regeneration. *Curr Opin Genet Dev.* 2003 Oct;13(5):497-501. PMID 14550415
2. Nye HL, Cameron JA, Chernoff EA, Stocum DL. Regeneration of the urodele limb: a review. *Dev Dyn.* 2003 Feb;226(2):280-94. PMID 12557206
3. Yu H, Mohan S, Masinde GL, Baylink DJ. Mapping the dominant wound healing and soft tissue regeneration QTL in MRL x CAST. *Mamm Genome.* 2005 Dec;16(12):918-24. PMID 16341671
4. Gardiner DM, Blumberg B, Komine Y, Bryant SV. Regulation of HoxA expression in developing and regenerating axolotl limbs. *Development.* 1995 Jun;121(6):1731-41. PMID 7600989
5. Torok MA, Gardiner DM, Shubin NH, Bryant SV. Expression of HoxD genes in developing and regenerating axolotl limbs. *Dev Biol.* 1998 Aug 15;200(2):225-33. PMID 9705229
6. Putta S, Smith JJ, Walker JA, Rondet M, Weisrock DW, Monaghan J, Samuels AK, Kump K, King DC, Maness NJ, Habermann B, Tanaka E, Bryant SV, Gardiner DM, Parichy DM, Voss SR. From biomedicine to natural history research: EST resources for ambystomatid salamanders. *BMC Genomics.* 2004 Aug 13;5(1):54. PMID 15310388
7. Andrews, Wyatt (March 23, 2008). "Medicine's Cutting Edge: Re-Growing Organs" (<http://www.cbsnews.com/stories/2008/03/22/sunday/main3960219.shtml>) . *Sunday Morning* (CBS News). <http://www.cbsnews.com/stories/2008/03/22/sunday/main3960219.shtml>.

## External links

- Spallanzani's mouse: a model of restoration and regeneration ([http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list\\_uids=14594211](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=14594211))
- Mice that regrow hearts in the news ([http://www.worldhealth.net/news/it\\_s\\_a\\_miracle\\_-\\_mice\\_regrow\\_hearts/](http://www.worldhealth.net/news/it_s_a_miracle_-_mice_regrow_hearts/))



- DARPA Grant Supports Research Toward Realizing Tissue Regeneration (<http://www.upmc.com/Communications/NewsBureau/Research/Articles/DarpaGrant.htm>)
- The Geniuses Of Regeneration ([http://businessweek.com/magazine/content/04\\_21/b3884008\\_mz001.htm](http://businessweek.com/magazine/content/04_21/b3884008_mz001.htm)) in *BusinessWeek*, May 24, 2004
- UCI Limb Regeneration Lab (<http://regeneration.bio.uci.edu>)

Retrieved from "[http://en.wikipedia.org/wiki/Regeneration\\_\(biology\)](http://en.wikipedia.org/wiki/Regeneration_(biology))"

Categories: Developmental biology

---

- This page was last modified on 12 April 2011 at 11:54.
- Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. See Terms of Use for details.  
Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.